

Daniel McIntyre

From: Liz Martin on behalf of Frannie Hutchinson
Sent: Tuesday, November 19, 2013 9:52 AM
To: Daniel McIntyre; Faye W. Outlaw
Subject: FW: Fertilizer Ordinance
Attachments: City of Cape Coral Fertilizer Ordinance Support 033010.pdf; Sarasota article excerpt.doc; Technical Support Document for EPC Fertilizer Rule - FINAL (1).pdf

fyi

From: David Botto [mailto:dbotto1@cfl.rr.com]
Sent: Tuesday, November 19, 2013 9:09 AM
To: Frannie Hutchinson
Subject: Fertilizer Ordinance

FROM: David Botto, Marine Resources Council (MRC)

Dear Commissioner;

I'm sure you are all aware of the devastating effects of recent algae blooms on our great natural treasure, the Indian River Lagoon. Excessive algae has killed over 47,000 acres of the sea grass on which almost every living thing depends. One acre of healthy sea grass produces 10,000 fish a year and tens of thousands of invertebrates such as shrimp and crabs. Such a loss means 470 million fish that won't be in our waters as well as the loss of most things fun to catch and good to eat. It doesn't end there, many of our favorite ocean critters start out in the lagoon sea grass and migrate to the ocean as juveniles - including grouper, snapper and even spiny lobster. The loss of our lagoon fishery is an economic catastrophe. The full effects are already showing up - fish catches are down, crabbing is almost gone, dipping shrimp on the bridges is a waste of time. Manatees, dolphin and pelicans are dying - some from starvation. Lagoon side restaurants no longer have the customer pleasing numbers of sea birds in the air and on the nearby docks, no fish, no sea fowl. Many of our most important businesses beyond those directly engaged in supporting commercial and recreational lagoon activity will suffer. Property values will fall as the waterfront turns to lifeless slime. This is happening and we can reverse it.

Scientists are not yet sure what triggered the killer algae blooms in the lagoon but one fact is indisputable - **excess algae can not exist unless there are excess nutrients in the water to feed it.** The lagoon is receiving almost 50% more nutrients, nitrogen and phosphorus, than it can safely absorb and use. That 50% is what is killing our fishery. There are several sources of the nutrients that are entering the lagoon. A major point source was sewage treatment plant effluent. SWIM legislation in the 90's put an effective end to that on the lagoon. Plant mods were very costly but nearly eliminated that source. We have spent many more millions to reduce run off nutrient pollution with capital intensive projects such as retention areas and baffle boxes. Brevard County data in the FLDEP final Basin Management Action Plan report indicate that our capital expenditure for storm water nutrient removal is over \$1000 PER POUND and annual O&M of these projects is over \$33 PER POUND.

The **Florida Consumer Fertilizer Task Force**, comprised of leaders from diverse backgrounds, firmly established in their 2004 final report that **urban turf fertilizer run off is a significant contributor to our nutrient pollution problem.** The Task Force recommended a state statute to manage the historical misuse and abuse of urban fertilizer. Unfortunately, special interest influence on its preparation so

weakened the state ordinance that it is ineffective and nearly impossible to follow. If you read it with the perspective of Joe Homeowner, its failings will be obvious. However, it does present a model and gives local government the authority to strengthen and clarify it to meet local needs. Over 50 Florida communities have done just that with great success. A strong fertilizer ordinance is proven to reduce nutrient pollution by 28% (Lehman 2009) and in some cases even more. The state ordinance has been in effect for over six years but the only waters in the state that have shown measured improvement in water quality are areas where stronger ordinances have been adopted. Tampa Bay National Estuary Program reports their sea grass has expanded at a rate of over 600 acres per year and they now have 1950's era grass coverage. In addition, lawn maintenance professionals in the Tampa Bay area report that yard grass is healthier and costs less to maintain. Excess fertilizing promotes fungus, chinch bugs and other diseases. Most of Florida's southwest coast communities have enacted strong ordinances and, without exception, they report no downside. Agribusinesses are thriving, lawns are healthier and cost less, enforcement is not a problem. The Florida Department of Environmental Protection has stated, in writing, that they will not challenge more stringent local fertilizer requirements. Most important, the cost to the taxpayer to help achieve a thriving fishery in this way is almost nothing.

There are some who claim that field lab tests done by agricultural scientists prove that healthy St. Augustine grass prevents nutrient pollution in leached, not run off, water. Others believe that the referenced tests were somewhat contrived for desired results and not independently peer reviewed. Both claims are irrelevant because your ordinance is concerned with surface run-off only and does not address leached water. It does not matter whether the science is good or bad because it does not apply to the ordinance being considered here. Incidentally, as mentioned earlier, smarter fertilizing has been shown to result in healthier yard grass while it's helping to restore the sea grass.

I urge you to enact a strong, effective fertilizer ordinance that will help put our lagoon fishery on the path back to a thriving economic engine for all our benefit. More years of study can not be allowed to stand in the way of this simple, almost no cost remedy. The lagoon fishery can't wait. I have attached studies and information for your review.

Respectfully,
David Botto, Chairman
Fisheries Campaign
TEL 321 773 0730

Please Note: Florida has very broad public records laws. Most written communications to or from County officials regarding County business are public records available to the public and media upon request. It is the policy of St. Lucie County that all County records shall be open for personal inspection, examination and / or copying. Your e-mail communications will be subject to public disclosure unless an exemption applies to the communication. If you received this email in error, please notify the sender by reply e-mail and delete all materials from all computers.

CAPE CORAL NORTH SPREADER CANAL ECOSYSTEM MANAGEMENT AGREEMENT

BACKGROUND INFORMATION IN SUPPORT OF PROPOSED LOCAL FERTILIZER ORDINANCE

December 11, 2009

Revised February 10, 2010

INTRODUCTION

Nutrients (nitrogen – N and phosphorus – P) have repeatedly been demonstrated to cause water pollution. Studies for the Charlotte Harbor National Estuary Program, Tampa Bay Estuary Program, and the Chesapeake Bay Program among many others have shown the adverse impacts of over-enrichment to surface waters. Fertilizer contains significant amounts of nutrients and must be considered in any program intended to protect and enhance surface water quality. The following summarizes a review of information supporting these provisions. Findings are taken from reports, studies, personal interviews, and computer modeling simulations.

The objective of this document is to provide technical assistance for the Cape Coral North Spreader Canal - Ecosystem Management Agreement (NSC-EMA) process. Specifically, information is summarized herein to augment the technical rationale for elements of the City of Cape Coral's (City) proposed ordinance regulating the application and use of landscape fertilizer (ordinance).

Section 373.62, Florida Statutes states that a local government may adopt additional or more stringent standards than the model ordinance if the following criteria are met:

- The local government has demonstrated, as part of a comprehensive program to address nonpoint sources of nutrient pollution which is science-based, and economically and technically feasible, that additional or more stringent standards to the model ordinance are necessary to adequately address urban fertilizer contributions to nonpoint source nutrient loading to a water body.
- The local government has documented its consideration of all relevant scientific information including input from the Florida Department of Environmental Protection (FDEP) and the Department of Agriculture and Consumer Services, if provided, on the need for additional or more stringent standards to address fertilizer use as a contributor to water quality degradation. All documentation must become part of the public record before adopting the additional or more stringent standards.

Some elements of the City's proposed fertilizer ordinance could be interpreted as being more stringent than the model ordinance developed by the FDEP. The following City ordinance provisions that are in this category are discussed below:

- Blackout period - no fertilizer application during the rainy season (June through September)
- Limited application rate – reduced from levels allowed in the State model ordinance
- Slow-release fertilizers – require the use of slow-release fertilizer
- No fertilizer shall be applied within ten (10) feet from any water body or seawall other than by careful hand dispersion methods which assure that no fertilizer is dispersed into the water.

The following discussion of these three provisions presents the scientific justification and potential cost ramifications of adopting an ordinance that includes each provision.

BLACKOUT PERIOD

Section 9-107 A. of the proposed ordinance states:

"No person shall apply fertilizers containing nitrogen and/or phosphorus to Turf and/or landscape Plants during the rainy season (June 1 through September 30 of each calendar year)."

Scientific Justification

The rationale for this provision is that more frequent rains during the summer will promote higher levels of runoff and leachate carrying nutrients to surface water and groundwater. This is reasonable, in that over 60% of our annual rains occur during the June through September period. Also, the brief but intense rains that are frequent (often daily) in the summer are ideal for suspending small particles such as fertilizer, dissolving it into its components, and carrying it to nearby water bodies. The University of Florida Institute of Food and Agricultural Sciences (IFAS) has recognized that stormwater runoff is a critical pathway for fertilizer transport to a receiving water body. Hauxwell et al. (2001) stated "Limit the use of fertilizers on residential and commercial lawns and landscaping. When fertilizing, it is important to minimize quantities and avoid fertilizing before heavy rains. Storm water runoff can transport fertilizer intended for yards directly to coastal waters." The Southwest Florida Water Management District (2009) also recommended "Do not fertilize if rain is predicted in the next 24–36 hours".

Hochmuth et al., (2009) concluded that summer application of fertilizer is beneficial because the warmer months are the peak growing season for turf grasses, and nutrient assimilation through

the roots is at a maximum. However, it is not necessary to facilitate maximum growth to achieve a healthy, attractive lawn. Moderate or low levels of growth maintain plant health and also reduce maintenance requirements. Allowing turf grasses and landscape plants to grow at a sustainable but less than maximum rate is a prudent, reasonable, and cost-effective approach to landscaping. This approach has been accepted by many. Lawn care specialists have observed that turf grasses not fertilized during the summer remain perfectly healthy (although sometimes not so emerald green during the summer months) but suffer no ill effects (Juchnowicz, 2009). Also, summer fertilization leads to more frequent mowing which stresses the grass and makes it more susceptible to damage from disease and insects (Juchnowicz, personal communication; Barth, personal communication).

Summer is not the only growth period for turf grasses and landscape vegetation. Plants are also active in spring and fall. SWFWMD (2009) recognized that the prudent application of fertilizer bracketing the rainy season is a reasonable alternative to summer fertilization, stating: *"Do not apply fertilizer if heavy rainfall is forecast in the next 24 hours."* and *"Fertilize only when the grass is actively growing. Spring and fall are the two key times for fertilizer application in Florida."*

In many cases, there are other sources of nutrients available for plants and turf. Reclaimed water is used for irrigation in many urban areas including Cape Coral, and has significant N content. Irrigation of turf using reclaimed water can greatly reduce the need for fertilizer application. On a large multi-use development in Sarasota County, changes to fertilization practices, including the use of reclaimed water, on turf grasses in common areas have reduced maintenance and turf repair costs by an order of magnitude and reduced required fertilizer rates to 1 lb/1000 ft²/yr or less (Barth, personal communication). Water management districts have long advocated the use of reclaimed water for irrigation, both to reduce water use and to lower nutrient application to lawns. For example, SWFWMD (2009) recommends *"If your household utilizes reclaimed water, you may not need to fertilize as much or as often. Reclaimed water contains nutrients like nitrogen and phosphorus, which are also in fertilizer"*. Usable nitrogen is also delivered to the land surface via atmospheric deposition.

There are other factors that should discourage summer applications of fertilizer. High N application rates have frequently been linked to increased problems with diseases such as fungus and insect pests such as chinch bugs. (Barth, personal communication) Summer is a peak growth season for insects and fungus as well as plants. Also, high N application during peak growing season can lead to "over-thatching" in St. Augustine grass (Barth, personal communication). This occurs when the thatch builds up to a depth where the grass root system cannot reach the soil, and obtains water and food only from the underlying grass layer. This makes the lawn more susceptible to drought, disease, and insect damage. Problems with

summer fertilization have also been recognized by IFAS "...fertilization with N in the summer is not always desirable since this often encourages disease and insect problems. Many times the addition of iron (Fe) to these grasses provides the desirable dark green color, but does not stimulate excessive grass growth which follows N fertilization" (Sartain, 2007).

Revised IFAS recommendations for turf grasses fertilization (Sartain, 2007) were examined. As noted, these recommendations focus solely on turf grass health and not the protection of water quality (Sartain et al, 2008). Many of the fertilization schedules recommend June or September applications. The potential impacts of shifting those applications one month should be investigated, and would likely cause no adverse effects. Also, addition of micronutrients such as iron during the summer is recommended in lieu of N in some instances, and might well be applicable for most or all situations.

If thoughtful practices are followed including using a slow-release fertilizer for the spring application, application of micro-nutrients such as iron during the summer, and not over-fertilizing in the spring or fall, no ill effects of the summer blackout should be observed. On the contrary, environmental benefits and cost savings will follow these practices.

In summary, fertilization during the summer may well stimulate turf grasses to their maximum growth rate. However, less robust growth during the summer will sustain an attractive, healthy lawn and greatly reduce the risk of pollution of surface waters and groundwater.

Cost Ramifications

There are potential cost savings that may be realized by not fertilizing during the summer. These include:

- reduced maintenance and labor cost resulting from slower growth rates and less frequent mowing,
- reduced cost for turf repair resulting from disease and insect damage, and
- less water required for irrigation.

Other cost benefits can also accrue, as discussed below.

SLOW-RELEASE FERTILIZERS

Section 9-107 B.2. of the proposed ordinance states:

"Fertilizers applied to Turf and/or Landscape Plants within City of Cape Coral shall contain no less than fifty percent (50%) Slow-release Nitrogen per Guaranteed Analysis Label."

Scientific Justification

The rate of absorption of slow-release fertilizer into soil and plants is slower than traditional fertilizer. This allows a more even and complete uptake of nutrients and reduces the risk of leaching into surface water and groundwater. A stated disadvantage of inorganic fertilizers is that they "leach readily" (Sartain, 2007; Zone Ten Nursery, 2009). Slow-release N helps to equalize the release rate of nutrients and lengthens the time period of effectiveness of an application, thus reducing the number of applications required over a year.

A slow-release fertilizer is safer to use than a highly soluble granular type as it reduces the risk of over-fertilization and burning plants. It also provides for proper levels of soil nutrients for a longer time. Fertilizers with only soluble N and P disappear rapidly from the soil.

Thus, the use of slow-release fertilizer presents an opportunity for a reduction in nutrients released into water bodies. These benefits have been extensively documented by commercial, academic, and regulatory entities (Blaylock et al., 2005; Wang and Alva, 1996; Zvomuya et al., 2003). Seedland (2009) stated *"The time-released fertilizers are easy for the homeowner to manage and won't burn the grass if sufficient rainfall occurs or systematic watering is available"*. The South Florida Water Management District (2009) also points out that slow-release fertilizer is an advantage when nutrients cannot be applied as frequently.

Many commercial groups understand the environmental benefits and economic opportunities of slow-release fertilizer. Georgia Pacific (2009) advertises that their Nitamin® brand fertilizers contain 60 – 94% slow-release N which:

- allows less ammonia volatilization,
- reduces nitrate leaching, and
- lowers required nitrogen application rates

Success with slow-release fertilizer has been demonstrated with commercial applications as well as research (American Vegetable Grower, 2009). Slow-release fertilizers come in many forms. Studies with ureaform (UF) products show beneficial effects of slow-release nitrogen fertilizer in improved fertility management and reduced pollution of drainage waters (Alexander and Ulrich Helm, 1990).

Thus, the proven technology of slow-release fertilizer provides less risk of environmental degradation and increased flexibility for the user. The extensive use of slow-release fertilizer is a valuable element of a comprehensive fertilizer management program.

Cost Ramifications

Cost savings that can accompany the use of slow-release fertilizer include the following:

- reduced labor cost due to fewer applications per year,
- reduced cost of repair of over-fertilized, burned turf, and
- less water required for irrigation.

Slow-release fertilizers are readily available and their cost is similar to traditional soluble product. Other cost benefits can also accrue, as discussed below.

LIMITED APPLICATION RATE

Section 9-107 B.3. of the proposed ordinance states:

"Fertilizers should be applied to Turf and/or Landscape Plants at the lowest rate necessary. No more than four (4) lbs. of nitrogen per 1000 ft² shall be applied to any Turf/landscape area in any calendar year."

Scientific Justification

Fertilization rates of N vary by region, plant type, weather, soil type, land use, and available level of maintenance. This is illustrated in IFAS recommendations for turf grass fertilization. Different application rates are listed for north, central, and south Florida, for different types of turf grass (St. Augustine grass, Bermuda grass, Bahia grass, etc.), and for different intensities of maintenance (low, medium, high) (Sartain, 2007).

The IFAS recommendations are not inconsistent with the proposed ordinance provision to limit application of N fertilizer to 4 pounds per 1000 square feet per year (lb/1000 ft²/yr). This value (4 lb/1000 ft²/yr) is within the range of, or even higher than, most of the recommended rates for south Florida turf grasses, so maintaining dosage below the "high" maintenance level will stay within the ordinance criterion. Also, the addition of micronutrients such as iron during the

summer is recommended in lieu of N in some instances, and might be applicable for most or all situations.

Finally, in determining the best application rate, all potential sources of N should be considered. As stated above, reclaimed water provides substantial N for plants. Assuming an N concentration of 10 mg/L in the water (typical for secondary treatment of wastewater) and an application rate of one inch per week for the 36-week dry season, a total of 1.3 lb/1000 ft²/yr of N would be delivered to the grass through the use of reclaimed water alone. This is a conservative estimate, as some irrigation would likely take place during the summer.

Limiting the amount of N applied to turf grasses seems to be a reasonable action. Providing adequate nutrients to sustain healthy grass can be achieved without high N inputs through fertilizer.

Cost Ramifications

Several cost savings can result from limiting application rates, including the following:

- reduced purchase cost as less product would be applied,
- reduced maintenance and labor cost resulting from slower growth rates and less frequent mowing,
- reduced cost for turf repair resulting from disease and insect damage, and
- less water required for irrigation.

Other cost benefits can also accrue, as discussed below.

BUFFER ZONE

It is proposed to include the following language in the City's fertilizer ordinance:

"No fertilizer shall be applied within ten (10) feet from any water body or seawall other than by careful hand dispersion methods which assure that no fertilizer is dispersed into the water."

Scientific Justification

Requiring a "no development" buffer zone along water bodies is a standard approach to protecting water quality from impacts due to nutrient enrichment, introduction of oil, grease, and heavy metals, and sediment (Correll, 1996). Numerous agencies in Florida now require a setback for site alteration and impervious surfaces, and protection of valuable habitat, within

buffer zones of 10 – 50 feet. Lee County requires a 50-foot buffer, except on Pine Island where the setback is greater (Lee County, 2009). The purpose of the buffer zones is to protect habitat, stabilize shorelines and reduce erosion, and limit pollutant (including nutrients) loading to the water bodies.

Keeping impervious surface away from shorelines reduces the potential for erosion and sedimentation in a water body. Likewise, not applying fertilizer adjacent to surface waters or wetlands greatly decreases the potential for over-enrichment of the water body and subsequent eutrophication. Numerous local governments in Florida now require a setback for site alteration and impervious surfaces, and protection of valuable habitat, within buffer zones of 10 – 50 feet (Environmental Protection Commission of Hillsborough County, 2007).

It is widely recognized by professional landscapers and researchers alike that maintaining a non-fertilized buffer strip along water bodies also is a good practice for protecting water. Buffer strips help absorb nutrients present in runoff as well as add a margin for application error (Watschke, 2010; White et al., 2007). Thus, setting a buffer for no or limited fertilization from water bodies and wetlands is well justified and documented.

Cost Ramifications

Several cost savings can result from restricting fertilization within a buffer adjacent to a surface water body or wetland including the following:

- reduced purchase cost as less product would be applied, and
- less need to improve degraded water quality through use of Best Management Practices

OTHER CONSIDERATIONS

The potential benefits of fertilizer management were illustrated in a recent study by the Tampa Bay Estuary Program (2008). Estimates of the potential reduction in N loadings to Tampa Bay were made for ranges of both percent contribution of fertilizer to total urban runoff nutrient loadings, and percent compliance with a series of recommendations for controlling fertilizer application, including a summer blackout with a possible one-time exemption. The effect of the most conservative values (percent contribution of 10% and compliance rate of 10%) yielded a reduction of only 0.3% in total N loading, but a percent contribution of 79% and compliance rate of 75% resulted in a 19.6% reduction in total N loading. Both the high end and low end of the ranges of values are likely extreme, so a reduction in N loading between 0.3% and 19.6% could be expected as a result of adoption of the City's ordinance.

There are good reasons to adopt an ordinance more stringent than the State's example. Among the most compelling is the fact that several water bodies within the City including Gator Slough, Horseshoe Canal, and Hermosa Canal, have been declared "impaired waters" with nutrients the causative agents by FDEP. Impairment is due to nutrient loadings to these water bodies exceeding that which can be assimilated in the water body while maintaining water quality standards and supporting a viable ecosystem.

Although there are many sources of nutrients that enter surface water and groundwater, it logically follows that, of the thousands of pounds of fertilizer that are put on the ground each year, enough enters the receiving water bodies to impact water quality and the local biota. Therefore, actions that reduce the volume of fertilizer applied, and to maximize the efficiency of uptake of the nutrients by plants, will benefit the environment.

In addition, more efficient fertilizer use will lead to cost savings through:

- Less fertilizer purchased
- Less time and effort in application
- Less mowing and other maintenance required
- Less repair of turf grasses due to fungus and insect damage
- Less water required for irrigation
- Less need to improve degraded water quality through use of Best Management Practices

The cost of cleaning polluted water is substantial, and local governments usually pay the majority of the amount. The ordinance provisions discussed above are a form of source control, that is, it reduces the amount of a pollutant before it enters the environment. This is a much more economical approach to protecting water quality than remediation, or treating the water after it has become polluted.

As part of the North Spreader Canal Ecosystem Management Agreement process, a conceptual design for a stormwater treatment area (STA) for the Gator Slough Basin was developed. Gator Slough is one of local water bodies determined to be impaired by FDEP, and the STA was designed to remove nitrogen from stream flow to address the impairment. It was determined that a site of at least 50 acres would be needed to provide adequate levels of treatment (to remove about 1,000 lb N per year).

Cost estimates for N removal from stormwater typically range from \$55/lb N removed (Coastal Environmental, 1998) to \$100/lb N removed (US EPA, 1999). Over a 25-year life of the STA approximately 25,000 lb N would be removed, so the construction cost could easily exceed

\$1,375,000. This does not include land purchase and operation/maintenance costs. The above cost estimate is consistent with estimates recently provided by the Tampa Bay Estuary Program of \$20 and \$100 per pound of N removed from water (Tampa Tribune, 2009).

An estimate can be made of potential cost savings from the above estimate that can be realized by implementing the City ordinance. The Gator Slough basin has approximately 10,500 acres, and an estimate can be made that 3,000 acres of the basin are fertilized. IFAS currently recommends using a total of 4.5 lb N/1,000 ft²/yr for medium maintenance St. Augustine grass in south Florida. If that amount is reduced by 0.5 lb N/1000 ft²/yr, a total of 65,340 lb N/yr will be used. The equivalent cost for removing that amount from water, at \$55/lb N, is \$3,593,000/yr. Even if only 5% of the 0.5 lb N/1000 ft²/yr reaches a water body, that still equals 81,675 lb or \$4,492,000 over the 25-year life of an STA.

Limiting a lawn growth rate will lower required levels of time and effort for maintenance, as well as provide a cost savings, as recognized by both academic and commercial publications. Trenholm et al. (2009) concluded *"A lower-fertility lawn is best for those with little time to spend on lawn care.... (Higher fertility lawns) ...will require more time and money for lawn care."* Seedland (2009) also stated that *"The level of maintenance you wish to provide (more fertilizer = more work, more mowing more disease control monitoring, and more water) will determine how many applications you make to your grass."*

Both environmental stewardship and cost savings can be realized while protecting the health and appearance of turf grasses and landscape vegetation by examining established traditional practices and adjusting them to reflect current priorities. The City's proposed fertilizer ordinance, with adequate enforcement, will be an effective tool to achieve these goals.

SUMMARY

Similar to most highly technical and developing fields, opinions about appropriate fertilizer management strategies differ among various entities. However, the discussion above presents credible, technically-defensible evidence that a suite of **science-based, and economically and technically feasible actions** can and should be implemented by the City. The turf and landscape industry has shown the ability to adapt to the adoption of similar ordinances in other jurisdictions.

Implementation of a source control approach to water quality protection is much less costly than cleaning the water after it has become polluted. It has been shown that keeping excess

fertilizer out of the environment is a superior method of environmental protection in terms of cost to local governments, citizens, and other land owners.

The provisions of the City's proposed fertilizer ordinance, including a summer blackout period, increased use of slow-release fertilizer, and promoting lower application rates, are necessary as part of an **effective, comprehensive approach to controlling nonpoint sources of pollution for the purpose of protecting and improving the City's aquatic and estuarine resources.**

REFERENCES

- Alexander, A. and H. Ulrich Helm. 1990. Ureaform as a Slow-release Fertilizer: A Review. In: Zeitschrift für Pflanzenernährung und Bodenkunde. V:153, n:4, pp: 249-255.
- American Vegetable Grower. March, 2009. Deliver Better Tubers.
<http://www.growingproduce.com/americanvegetablegrower/index.php?storyid=1462>
- Barth, R. 2009. President, Palmer Ranch Master Property Owners Assoc. Personal Communication.
- Blaylock, A.D., J. Kaufmann, and R.D. Dowbenko. 2005. Nitrogen Fertilizer Technologies. In: Proceedings, Western Nutrient Management Conference. 2005. Vol. 6. Salt Lake City, UT.
- Cape Coral, City of. 2009. An Ordinance Amending Chapter 9 of the City Code Creating Article VI - Sections 9-100 to 9 - 13; Providing for the Application and Use of Fertilizers - draft. Cape Coral, FL
- Coastal Environmental/PBS&J, Inc. 1998. The Lake Tarpon Basin Management Plan. Prepared for Pinellas County. Clearwater, FL.
- Correll, D.L. 1996. Buffer zones and water quality protection: general principle. In: Buffer Zones: Their Processes and Potential in Water Protection. The Proceedings of the International Conference on Buffer Zones September 1996. Edited by N.E. Haycock, T.P. Burt, K.W.T. Goulding, and G. Pinay. Quest Environmental. P.O. Box 45, Harpenden, Hertfordshire, AL5 5UJ, UK.
- Environmental Protection Commission of Hillsborough County. 2007. EPC staff technical findings and recommendations in response to CARE Policy 19.1 of the Comprehensive Plan. Tampa, FL.
- Florida Fertilizer Task Force. 2007. Comments on draft State model ordinance. Tallahassee, FL.
- Georgia Pacific. 2009. Nitamin brand fertilizer specifications.
<http://www.gp.com/Plant Nutrition/Default.aspx>
- Greenview Summer. 2008. Fertilizing: when, why and how.
<http://www.greenviewblog.com/2008/04/fertilizing-y-2.html>
- Hauxwell, J., C. Jacoby, T. K. Frazer, J. 2001. Nutrients and Florida's Coastal Waters: The Links Between People, Increased Nutrients, and Changes to Coastal Aquatic Systems. Publication SGEB-55. Florida Sea Grant College Program. Gainesville, FL.

Hochmuth, G., T. Nell, J. Sartain, B. Unruh, M. Dukes, C. Martinez, L. Trenholm, and J. Cisar. 2009. Unintended Consequences Associated With Certain Urban Turf and Landscape Ordinances. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.

Juchnowicz, M. 2009. President, Gardenmasters of Southwest Florida, Inc. Personal Communication.

Kimball, J.W. 1994. Kimball's Biology Pages – The Nitrogen Cycle.
<http://users.rcn.com/jkimball.ma.ultranet/BiologyPages/N/NitrogenCycle.html>

Lee County Board of County Commissioners. 2009. Land Development Code - Ordinance No. 09-23, Ft. Myers, FL.

Sartain, J.B. 2007. General Recommendations for Fertilization of Turfgrasses on Florida Soils. IFAS #SL21. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.

Sartain, J.B., L.T. Trenholm, E.F. Gilman, T.A. Obreza, and G. Toor. December 2008. Frequently Asked Questions about Landscape Fertilization for Florida-Friendly Landscaping Ordinances. IFAS #ENH1115. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.

Seedland Fertilizers. 2009. Welcome to Lawnfertilizers.com.
<http://www.lawnfertilizers.com>

South Florida Water Management District. 2009 Western Broward County Turf & Landscape Best Management Practices (BMPs).

Southwest Florida Water Management District. 2009. Fertilizer Facts. Brooksville, FL.
http://www.swfwmd.state.fl.us/yards/fertilizer_facts/details.php#whentoapply

Tampa Bay Estuary Program. 2008. Technical Memorandum: Model-Based Estimates of Nitrogen Load Reductions Associated with Fertilizer Ordinance Implementation. St. Petersburg, FL.

Tampa Tribune. 2009. Fertilizer ban will save tax dollars. Editorial December 2, 2009. Tampa, FL.

Trenholm, L.E., J.L. Cisar, and J. Bryan Unruh. 2006. St. Augustine grass for Florida Lawns. Publication ENH5 Institute of Food and Agricultural Sciences. University of Florida. Gainesville, FL.

Trenholm, L.E.. 2008. Urban Turf Fertilizer Rule for Home Lawn Fertilization. Publication ENH1089 Institute of Food and Agricultural Sciences. University of Florida. Gainesville, FL.

US EPA. 1999. Costs and Benefits of Storm Water BMPs. US Environmental Protection Agency, Office of Science and Technology. Washington, D.C.

Wang, F.L. and A.K. Alva. 1996. Leaching of Nitrogen from Slow-Release Urea Sources in Sandy Soils. *Soil Sci Soc Am J* 60:1454-1458

Watschke, T.L. Proper use of fertilizers minimizes environmental effects. *Grounds Maintenance for Golf and Green Industry Professionals*. 2010.
http://www.grounds-mag.com/mag/grounds_maintenance_proper_fertilizers_minimizes/

White, S.K., H. F. Cook, and L. Garraway. 2007. The Use of Fertilizer-Free Grass Buffer Strips to Attenuate Nitrate Input to Marshland Dykes. *Water and Environment Journal*. Volume 12 Issue 1, Pages 54 - 59

Zone Ten Nursery. 2009. Tree Planting. Miami, FL.
<http://www.zoneten.com/>

Zvomuya, F., Rosen, C.J., Russelle, M.P., and Gupta, S.C. 2003. Nitrate leaching and nitrogen recovery following application of polyolefin-coated urea to potato. *J. Environ. Qual.* 32:480-489.

**ENVIRONMENTAL PROTECTION COMMISSION
OF HILLSBOROUGH COUNTY**

**TECHNICAL SUPPORT DOCUMENT FOR PROPOSED LOCAL
FERTILIZER RULE – CHAPTER 1-15**

May 7, 2010

INTRODUCTION

In an effort to improve water quality in Hillsborough County, the Environmental Protection Commission of Hillsborough County (EPC) is preparing a draft fertilizer management rule which is stricter than the model code created by the Florida Department of Environmental Protection (FDEP).

Excessive inputs of the macro-nutrients nitrogen (N) and phosphorus (P) have been demonstrated to cause water pollution (see, for example, entire December 2008 edition of the journal *Harmful Algae* (2008) as well as Southwest Florida Water Management District (SWFWMD) 2002. Multiple studies by and for the Environmental Protection Commission of Hillsborough County, Tampa Bay Estuary Program (TBEP), and the Chesapeake Bay Program among others have shown the adverse impacts of over-enrichment to surface waters. A recent scientific study by Peebles et al. (2009) for an impaired watershed partially contained within Hillsborough County indicates that a significant and growing percentage of the nitrogen in sediments is coming from inorganic fertilizer sources. Leggette, Brashears & Graham, Inc. (LBG) (2004) examined sources of nitrogen to groundwater in Lake Tarpon, a large lake near the Hillsborough County/Pinellas County border. They estimated that roughly 79% of the nitrogen in the surficial groundwater system contributing to Lake Tarpon is from fertilizers (agricultural and residential). Most fertilizers contain significant amounts of nutrients and must be considered in any program intended to protect and enhance surface water quality. The following summarizes a review of information supporting these provisions. These findings are taken from reports, studies, personal interviews, and public workshop comments.

The objective of this document is to provide technical assistance for the Commissioners, FDEP, Florida Department of Agriculture and Consumer Services (FDACS), and the University of Florida's Institute of Food and Agricultural Sciences (IFAS). Specifically, information is summarized herein to augment the technical rationale for elements of EPC's proposed rule regulating the application, purchase, and use of landscape fertilizer within Hillsborough County.

Section 403.9337, Florida Statutes states that a local government may adopt additional or more stringent standards than the model ordinance if the following criteria in subsection (2)(a) and (b) are met:

403.9337 Model Ordinance for Florida-Friendly Fertilizer Use on Urban Landscapes.--

(1) All county and municipal governments are encouraged to adopt and enforce the Model Ordinance for Florida-Friendly Fertilizer Use on Urban Landscapes or an equivalent requirement as a mechanism for protecting local surface and groundwater quality.

(2) Each county and municipal government located within the watershed of a water body or water segment that is listed as impaired by nutrients pursuant to s. 403.067, shall, at a minimum, adopt the department's Model Ordinance for Florida-Friendly Fertilizer Use on Urban Landscapes. A local government may adopt additional or more stringent standards than the model ordinance if the following criteria are met:

(a) The local government has demonstrated, as part of a comprehensive program to address nonpoint sources of nutrient pollution which is science-based, and economically and technically feasible, that additional or more stringent standards than the model ordinance are necessary in order to adequately address urban fertilizer contributions to nonpoint source nutrient loading to a water body.

(b) The local government documents that it has considered all relevant scientific information, including input from the department, the institute, the Department of Agriculture and Consumer Services, and the University of Florida Institute of Food and Agricultural Sciences, if provided, on the need for additional or more stringent provisions to address fertilizer use as a contributor to water quality degradation. All documentation must become part of the public record before adoption of the additional or more stringent criteria.

Some elements of the EPC's proposed fertilizer rule are more stringent than the model ordinance developed by the FDEP. The following provisions that are in this category are discussed below:

- Seasonal Restrictions - no N or P fertilizer application during the rainy season (June 1 - September 30)
- Retail Sales Restriction - no N or P fertilizer sales during restricted season (June 1- September 30)
- Fertilizer-Free Zones - No fertilizer shall be applied within ten (10) feet from any surface water body or seawall.

This document is the EPC's demonstration and justification for a more stringent fertilizer rule, but this condensed paper does not reflect the entirety of the documents the EPC has reviewed during this process. Nonetheless, this paper cites the most relevant research to support the EPC's position.

NONPOINT SOURCE POLLUTION PROTECTION

The EPC is an independent local government and separate from the government of Hillsborough County. Hillsborough County also has three municipalities: City of Plant City, City of Tampa, and City of Temple Terrace. The County and the three municipalities all have waterbodies impaired for nutrients, thus pursuant to section 403.9337, F.S. those four governments must adopt fertilizer management ordinances. The EPC, in coordination with Hillsborough County and the three municipalities, is proposing a draft fertilizer rule that will apply within all four of these governments' jurisdictions. The FDEP has concurred with the EPC that the EPC can pass one rule to fulfill all four governments' obligations under section 403.9337, F.S.

The four local governments and the EPC have comprehensive programs in place to address nonpoint sources of nutrient pollution. The comprehensive programs have been effective in improving surface water quality and seagrass restoration in the Tampa Bay region, but additional and more stringent fertilizer regulations are needed than the FDEP's model ordinance in order to adequately address urban fertilizer contributions to nonpoint source nutrient loading.

In fact, through the Hillsborough River Basin Management Action Plan (FDEP, 2009), the Tampa Bay region addresses surface water quality improvement of these impacted waters. Area stakeholders have provided resources of more than \$80 million dollars for operational and program costs to improve surface water quality in this area (see BMAP attachment).

Additional examples of comprehensive programs to address nonpoint source nutrient pollution include:

Wastewater

- Advanced Wastewater Treatment Levels on all Plants discharging to Tampa Bay and tributaries
- Industrial Pre-treatment requirements
- Private Pump Station Identification and Compliance Program
- Emergency Response Program
- Sanitary Sewer Overflow Database
- Contamination Complaint Response Program
- Lift Station Security Program
- Septic Tank Mapping and Hot Spot Program
- Septic Tank Set-back Program (Land Development Code)
- Public Health Education Program
- Environmental Protection Commission Regulatory Authority
- EPC water quality monitoring program

Regional and Miscellaneous Efforts

Watering Restrictions
 Maximization of Use of Reclaimed Water on Lawns
 Promotion of Florida Friendly Yards and Neighborhoods
 Cooperative Extension Education Programs
 Best Management Action Plans Adopted for multiple impaired water bodies
 Tampa Bay Estuary Program
 Agency for Bay Management/Tampa Bay Regional Planning Council
 Area-wide Water Quality Plan
 SWFWMD's Surface Water Improvement and Management Program
 Hillsborough River Watershed Alliance
 Tampa Bay Comprehensive Conservation and Management Plan (CCMP)
 Nitrogen Management Consortium
 Development of Multiple Watershed Management Plans
 Reasonable Assurance document
 Municipal NPDES stormwater permits
 EPC MOUs with Temple Terrace, County, and Tampa to assist in compliance and enforcement of their NPDES permits related to stormwater and illicit discharges
 Cities and County costly infrastructure upgrades to improve stormwater treatment
 TBEP, Tampa Bay Water, Hillsborough, Tampa, and SWFWMD public messaging regarding fertilizer use, water conservation, Florida-friendly planting, and/or irrigation practices.

Stormwater

County Stormwater Quality Management Ordinance #94-15
 City of Temple Terrace Code of Ordinances, Section 25.716
 Stormwater Ecologist Education Program
 Pet Waste Campaign Study
 Lake Watch and Hillsborough County Stream Water Watch
 Adopt-a-Pond Program
 Systematic Program for Stormwater Improvements
 Separate Stormwater Utility Fee
 Conservation Element of the County Comprehensive Plan
 Stormwater Management Element of the Comprehensive Plan
 Street sweeping that the County and Tampa participate in through FDOT
 EPC water quality monitoring program
 Officer Snook Education Program

Clearly, the Tampa Bay region is committed to reducing nonpoint source pollution, but best on impaired waters in the region and water quality data showing continued nutrient pollution, a more stringent rule is merited.

SEASONAL RESTRICTIONS

Section 1-15.04(a) of the proposed rule states:

"No applicator shall apply fertilizers containing nitrogen and/or phosphorus to turf and/or landscape plants during the restricted season from June 1 -September 30.

Justification

National Weather Service data (NOAA 2010 and WeatherBase.com 2010) for the Tampa Bay region indicates that the months of June, July, August, and September are the 4 wettest months of the year. Randy Lascody, Senior Forecaster, of the National Weather Service (NOAA 2002) describes the climatological reasons for this. In addition to tropical storms and hurricanes, these 4 months are dominated by daily thunderstorm activity in the Tampa Bay region. The exact locations of the rainfall associated with these thunderstorms cannot be easily predicted by meteorologists, nor can the intensity of a given rain event. The state fertilizer model does not contain a summer application ban, but in some regions of the state the summer rains are not as frequent or intense as in the Tampa Bay region, thus the one-size fits all fertilizer model does not make sense for a region that experiences different rainfall patterns than certain other regions of Florida.

Rainfall data over the last 48 years for Tampa clearly demonstrates that the preponderance of yearly rain falls during the months of June, July, August and September (WeatherBase.com 2010). More specifically, over the past 48 years of record, June rainfall averaged 5.6", July 7.3", August 7.9", and September 6.3". Combined, these 4 months represent more than half the average annual total rainfall received. The next highest month was March, with 3.4". In addition to tropical storms and hurricanes, these 4 months are dominated by thunderstorm activity. Over the past 48 years of record, the average number of days with thunderstorm activity in Tampa has been as follows: 14 days in June, 21 days in July, 20 days in August, and 12 days in September. The next closest month is May, with an average of 5 days. Finally, over the past 48 years, on average June has had 12 rainy days, July - 16, August - 17, and September - 13. The next closest months are January, February, March, and October, with 7 rainy days on average each.

The rationale for this provision to ban the application of N and P containing fertilizers during this 4 month period (termed "summer period" for purposes of this report) is that more frequent rains will increase the likelihood of higher levels of soil saturation, runoff, and leachate carrying nutrients to surface water and groundwater. This is reasonable, in that over 50% of our annual rains occur during this summer period (Weatherbase.com 2010). Also, the brief but intense rains create the ideal conditions for

suspending small particles such as fertilizer, dissolving it into its components, and carrying it to nearby water bodies. IFAS has recognized that stormwater runoff is a critical pathway for fertilizer transport to a receiving water body. Hauxwell et al. (2001) states "limit the use of fertilizers on residential and commercial lawns and landscaping. When fertilizing, it is important to minimize quantities and avoid fertilizing before heavy rains. Storm water runoff can transport fertilizer intended for yards directly to coastal waters." SWFWMD (2009) also recommends "do not fertilize if rain is predicted in the next 24–36 hours." Since it is extremely difficult to predict whether a given location will receive rain on a given day during the summer period, having a general ban on the spreading of N and P fertilizer during these 4 months would provide assurances that excess nutrients are not being washed into surface water bodies. Additionally, given the high frequency of rainfall during the summer months (e.g. 21 days in July), it is difficult to refrain from fertilizing within 24 to 36 hours if it may rain every other day or more frequently in the summer.

According to IFAS staff, a prohibition on the application of N-fertilizer during the 4 months of frequent rains has the potential to "stress" some lawns and landscapes. Multiple comments received from representatives of the lawn/landscape industry as well as from representatives of fertilizer manufacturers and distributors also echo this sentiment. Hochmuth et al. (2009) concluded that summer application of fertilizer can be beneficial because the warmer months are the peak growing season for turf grasses, and nutrient assimilation through the roots is at a maximum. IFAS staff, as well as representatives from the landscaping industry, also express concerns that a 4 month time period may be too extreme and can result, over time, in lowered turf grass health, hence lowered assimilation capabilities. This view is echoed by representatives from FDACS (Rackley and Collier, 2010, personal communication) and FDEP (Thomas, 2010, personal communication), among others. Given there are already fertilizer products on store shelves that claim to be slow-release for time frames that are 4 months or longer, it would appear that such products could be applied prior to the application ban and still provide enough nutrition to maintain turf health for the entire period. FDACS, IFAS, and others suggest that such claims are not realistic under high summer temperatures and high moisture conditions (both of which are factors that break down the slow-release coatings on granular slow release fertilizer products). However, we have been able to find no long-term scientific study to definitively address the question of the efficacy of slow-release fertilizer products to last over a 4 month summer period. It is our position, at this time, to accept the claims made as they are stated on the labels of the fertilizer products. We welcome scientific data that suggest that these claims are not valid.

A recently-received study of N-deficiency in turf grasses from Sarasota County (which has a summer application ban) compared with lawns in neighboring counties without a summer ban appears to indicate a higher percentage of N-deficient lawns in Sarasota compared with those other neighboring counties (Troutman, et al., 2008). The authors suggest that the N-deficiency observed will tend to increase over the coming years of continued application ban. The study does not, however, address water quality run-off from the various lawns. In fact, the majority of turf grass studies performed in Florida have focused primarily on requirements for optimal turf grass health rather than on quantifying run-off of nutrients to surface waters during rain events.

One question that remains largely unanswered in the literature is whether it is necessary to facilitate maximum turf growth to achieve and maintain a healthy, attractive lawn. Some landscapers have provided their opinions to EPC staff that moderate or low levels of growth maintain plant health and also reduce maintenance requirements (public workshop comments). These landscapers suggest that allowing turf grasses and landscape plants to grow at a sustainable, but less than maximum rate is a prudent, reasonable, and cost-effective approach to landscaping. Some lawn care specialists have observed that turf grasses not fertilized during the summer remain perfectly healthy (although sometimes not so emerald green during the summer months), but suffer no ill effects (Juchnowicz, 2009).

For much of Florida, including the Tampa Bay region, summer is not the only growth period for turf grasses and landscape vegetation. Plants are also active in spring and fall. SWFWMD (2009) recognized that the prudent application of fertilizer bracketing the rainy season is a reasonable alternative to summer fertilization, stating: *"do not apply fertilizer if heavy rainfall is forecast in the next 24 hours."* and *"fertilize only when the grass is actively growing. Spring and fall are the two key times for fertilizer application in Florida."*

It is also necessary to mention that N from inorganic fertilizers is not the only source of N available to plants (or that may end up in surface waters) during summer months. In many cases, there are other sources of nutrients available for plants and turf. Reclaimed water containing N is used for irrigation in many areas including much of urbanized unincorporated Hillsborough County, parts of Tampa and Plant City. Irrigation of turf using reclaimed water can reduce the need for fertilizer application. Barth (2009, personal communication) provided his assessment of reclaimed water on a large multi-use development in Sarasota County. He indicated that changes to fertilization practices, including the use of reclaimed water, on turf grasses in common areas have the potential to reduce maintenance and turf repair costs by an order of magnitude. Since December 2006 their fertilizer usage rates were reduced to 4 lb/1000 ft²/yr

or less. However, Barth does not suggest a definitive application rate. Water Management Districts have long advocated the use of reclaimed water for irrigation, both to reduce water use and to lower nutrient application to lawns. For example, SWFWMD (2009) recommends *"if your household utilizes reclaimed water, you may not need to fertilize as much or as often. Reclaimed water contains nutrients like nitrogen and phosphorus, which are also in fertilizer."* Usable nitrogen is also delivered to the land surface via atmospheric deposition and from organic N decomposing in grass clippings, leaves, etc.

The debate relating to the summer application ban extends to the effects of N-fertilization on disease and pest infestations as well. High N application rates have been linked to increased problems with diseases such as fungus and insects such as chinch bugs (public workshop comments). Summer is a peak growth season for insects and fungus as well as plants. Problems with summer fertilization and disease/insect problems have also been recognized by IFAS *"..fertilization with N in the summer is not always desirable since this often encourages disease and insect problems. Many times the addition of iron (Fe) to these grasses provides the desirable dark green color, but does not stimulate excessive grass growth which follows N fertilization"* (Sartain, 2007). However, other persons familiar with the same areas of Sarasota County as used by Barth for his opinions have stated that these observations pre-dated the applications of new pesticides on these areas and that observations of reduced pests and fungus are actually the result of these new products, rather than attributable to reduced N applications (public workshop comments).

Barth (2010, personal communication) also indicated that high N application during peak growing season can lead to "over-thatching" in St. Augustine grass. This occurs when the thatch builds up to a depth where the grass root system cannot reach the soil, and obtains water and food only from the underlying grass layer. This makes the lawn more susceptible to drought, disease, and insect damage.

It should be noted that no landscape professional we have taken public comments from has advocated summer fertilization with high concentrations of N. The differences of professional opinions expressed at our public workshops have centered on whether it is more beneficial to allow "spoon-feeding" (limited applications of 0.5 lbs/1000 sq. ft. or less on turf and/or landscape plants during summer months). This would, according to some, help avoid turf stress versus the increased chances for N run-off into surface water bodies as a result of stressed lawns and frequent summer-time rain events.

Revised IFAS recommendations for turf grass fertilization (Sartain, 2007) were examined. As noted, these recommendations focus primarily on turf grass health and not directly on the protection of surface

water quality (Sartain et al. 2008). Many of the fertilization schedules recommend June or September applications. The potential impacts to turf grass conditions of shifting those applications one month should be investigated in controlled, field-based experiments. IFAS publications plus comments by multiple persons indicate that the addition of micronutrients such as iron during the summer is recommended with, and in some cases, in lieu of N. However, these same persons acknowledge that a small concentration of N is needed for efficient uptake of the micronutrients by the plant.

One potential approach, proposed by many landscaping industry representatives, to help minimize the misuse of fertilizer during the summer rainy season while maintaining some flexibility to attend to turf and/or landscape plants that may have a nitrogen deficiency would be to require that all persons who apply fertilizer during the summer months be trained and certified. We have been unable to locate data on the potential for pollution as a result of such an exemption. A study conducted in the Wekiva River watershed (FDEP/SJRWMD 2010) suggests that commercial landscape/lawn care companies do contribute substantial amounts of N to surface waters. However, this study did not differentiate between trained and untrained lawn care professionals.

Another area of uncertainty relates to landscape plants. There is general agreement that landscaped areas represent a small percentage of the typical yard and that "spoon feeding" of landscape plants represents a smaller chance of contributing to surface water run-off of nutrients than does broadcast-type fertilization of turf. There is NOT general agreement, however, over the *need* to fertilize landscape plants during the summer months. Information from some landscape plant nurseries suggests that newly planted landscape plants will require N inputs within a few months of initial planting, hence a 4 month blackout period may prove detrimental to the survival of newly planted landscape plants. Conversely, some nurseries sell their plants pre-fertilized thereby eliminating the need for additional nutrients upon initial placement in the ground.

Another aspect of a proposed application ban during the summer rainy season that is difficult to quantify is the cost/benefit ratio. Hillsborough County engineers and scientists within the stormwater and water resource services departments are currently examining the potential costs to reduce N concentrations in impaired water bodies. Current estimates run into tens, if not hundreds, of millions in additional tax dollars for infrastructure, operations, and maintenance costs (HCPW 2009, personal communication). Efforts to preempt the possible introduction of nutrients into surface waters during summer rainy periods appear to be a relatively less expensive tool to implement given the massive costs of non-compliance. Additionally, proponents of an application ban suggest that individual homeowners will save money via

less effort to cut turf due to slower growth rates with less N, less costs to treat for disease and pests, and less cost for irrigation. Opponents of fertilizer application and sales restrictions may argue that there is also the potential for over-use of fertilizer in May and October with a resultant excess of nutrients moving into surface water bodies. Others suggest that an outright ban on N and P fertilization for 4 months will reduce turf health over time, leading to increased costs for turf restoration plus increased use of N to bring the turf back to health.

Sarasota County has had a ban on N-fertilizer applications during the 4 month summer rainy season since 2008. Consequently, their application ban has been in place for 2 summer rainy seasons. Documentation from Sarasota County officials indicate that they have seen no overall decline in turf conditions (Sarasota County BOCC, 2009). However, at least one IFAS turf specialist who visited Sarasota during the fall of 2009 returned with an opinion that did not fully match that of the Sarasota officials (Hochmuth 2009, personal communication).

On balance, several points are evident with respect to the issue of the pros and cons of a summertime application ban:

1. It is reasonable to conclude that a summertime fertilizer ban will result in a decrease in nutrient inputs to surface water bodies during that time frame;
2. Overall costs of such a ban to the average Hillsborough County homeowner will likely be far less than the costs of subsidizing additional stormwater infrastructure and subsequent operation and maintenance of infrastructure to treat nutrient laden waters;
3. The ability of meteorologists to pin-point specific locations for summer rain and to predict expected quantities of rain from a given storm event remains problematic therefore, there is always a chance of nutrients from a morning fertilization becoming part of a surface water run-off event that afternoon or the following day.

However, multiple aspects of this issue remain debatable without additional data:

1. The extent to which the inability to use N/P for turf/landscape applications during 4 months might gradually reduce plant health such that the beneficial aspects of lawns and landscapes (sediment/nutrient filtering, CO₂ sequestration, thermal reductions, etc.) might be compromised and would then require additional N to restore over and above what would be required for maintenance;
2. The potential for negative economic impacts on the industry;
3. Whether turf with no summertime inorganic N is less or more resistant to pests and disease;
4. Whether allowing trained/licensed professionals to use fertilizers in the summer, as their training deems necessary, will increase the amount of N run-off into surface water bodies when compared with an absolute ban;
5. The effectiveness of slow release fertilizers to last for more than 4 months during a summertime ban.
6. Whether very low concentrations of N in fertilizers and/or top dressings that contain micro-nutrients would improve plant health without compromising water quality.

On balance, the weight of evidence with respect to an application ban during the summer rainy season suggests that it would be prudent and more cost-effective to minimize the opportunities for nutrient runoff into surface water bodies by prohibiting applications containing N/P. Manufacturers' labels indicate that there are products currently on the market that can last for more than the 4 month application ban period. Some evidence from 2 summers of bans in Sarasota indicates that turf health has not declined dramatically, if at all.

The Commission may want to closely examine an option that allows landscape plants to be "spoon-fed" during the summer ban. The potential for water quality degradation is less likely for selective use of N-fertilizers for landscape plants than with broadcast spreading for turf. The Commission may also choose to examine an option that would allow trained and licensed lawn care professionals only to apply N/P fertilizer during the summer ban period should the turf and/or landscape plants need a small concentration of N to maintain health. It should be noted again, however, that we have found no scientific documentation quantifying potential water quality improvements using this approach.

SLOW-RELEASE FERTILIZERS

Section 1-15.05 of the proposed rule requires a minimum percentage of slow-release granular nitrogen fertilizer.

Justification

The rate of absorption of slow-release, inorganic fertilizer into soil and plants is slower than traditional fertilizer. This allows a more even and complete uptake of nutrients and reduces the risk of leaching into surface water and groundwater. A stated disadvantage of typical, inorganic fertilizers is that they "leach readily" (Sartain, 2007; Zone Ten Nursery, 2009). Slow-release N helps to equalize the release rate of nutrients and lengthens the time period of effectiveness of an application, thus reducing the number of applications required over a year.

A slow-release fertilizer is safer to use than a highly soluble granular type as it reduces the risk of over-fertilization and burning plants. It also provides for proper levels of soil nutrients for a longer time. Fertilizers containing only soluble N and P disappear rapidly from the soil. Thus, the use of slow-release fertilizers present an opportunity for a reduction in nutrients released into water bodies. These benefits have been extensively documented by commercial, academic, and regulatory entities (Blaylock et al., 2005; Wang and Alva, 1996; Zvomuya et al., 2003). Seedland (2009) stated "*the time-released fertilizers are easy for the homeowner to manage and won't burn the grass if sufficient rainfall occurs or systematic*

watering is available." The South Florida Water Management District (2009) also points out that slow-release fertilizer is an advantage when nutrients cannot be applied as frequently.

Some commercial groups understand the environmental benefits and economic opportunities of slow-release fertilizer. Georgia Pacific (2009) advertises that their Nitamin® brand fertilizers contain 60 – 94% slow-release N which:

- allows less ammonia volatilization,
- reduces nitrate leaching, and
- lowers required nitrogen application rates

Success with slow-release fertilizer has been demonstrated with commercial applications as well as research projects (American Vegetable Grower, 2009). Slow-release fertilizers come in many forms. Studies with ureaform (UF) products show beneficial effects of slow-release nitrogen fertilizer in improved fertility management and reduced pollution of drainage waters (Alexander and Ulrich Helm, 1990).

Thus, the proven technology of slow-release fertilizer provides less risk of environmental degradation and increases flexibility for the user. The extensive use of slow-release fertilizer is a valuable element of a comprehensive fertilizer management program.

Cost / Benefit to Citizens

Cost savings that can accompany the use of slow-release fertilizer include the following:

- reduced labor cost due to fewer applications per year,
- reduced cost of repair of over-fertilized, burned turf, and
- less water required for irrigation.

Slow-release fertilizers are readily available in Florida, but their cost is currently higher than traditional soluble product.

The data indicate that the use of slow release fertilizers can be beneficial. The EPC endorses the use of slow-release fertilizers. The proposed rule follows the TBEP model ordinance and remains silent as to exact percentage(s) of slow-release N-fertilizer required.

RETAIL SALES RESTRICTION

Section 1-15.11(a) of the proposed rule states:

"Effective June 1, 2012, no person shall sell, at retail, any lawn or landscape fertilizer, liquid or granular, within Hillsborough County that contains any amount of nitrogen or phosphorous during the restricted season from June 1-September 30, unless otherwise provided for in this rule."

Justification

Decades of surface water quality monitoring data by the EPC clearly indicate the negative impact that the summer rainy season has on the already nutrient impaired waters of Tampa Bay. In an earlier section of this report we have presented the arguments for and against the summer time application of N-fertilizer. These same arguments apply to the sales restriction issue. From a regulatory/enforcement perspective, the prohibition of sales is more readily enforceable than is the prohibition of application.

It has been suggested by representatives of the landscaping industry, representatives of the fertilizer manufacturers, FDACS staff, FDEP staff, and IFAS staff that a summer sales restriction would have "unintended consequences" including, but not limited to:

- compromised turf grass health and viability
- increased stormwater runoff due to compromised lawn's inability to filter water
- over fertilization by homeowners immediately prior to and following the summer ban period leading to increased nutrient loading
- inconsistency in fertilizer regulations throughout Florida

However, it has also been suggested by various local governments and TBEP that limiting the summer sale of fertilizers would:

- prevent excessive nutrients from entering surface waters due to misuse of products and application techniques
- provide a consistent practice and message throughout the Tampa Bay area since Pinellas County has already passed a similar sales restriction
- allow local governments a cost-effective means of reducing additional nutrient loads to already impaired waters of Hillsborough County
- be a cost-effective mechanism of enforcement and education

Given the current lack of research on the subject, the most common sense approach would suggest that fewer nutrients placed on lawns and impervious surfaces during summer months would lead to fewer opportunities for nutrient runoff and, therefore, be beneficial to the surface waters of Hillsborough County particularly during those months of increased rainfall activity. Florida case law has not addressed fertilizer sales bans, but it has been addressed in federal court in Wisconsin and upheld. (see Croplife America Inc. et al vs. City of Madison, et al, 373 F.Supp.2d 905)

Cost / Benefit to Citizens

Several cost savings can result from a retail sales restriction, including the following:

- reduced annual purchase cost as less product would be purchased and applied,
- reduced maintenance and labor cost resulting from slower growth rates and less frequent mowing,
- reduced cost for turf repair resulting from disease and insect damage, and
- less water required for irrigation.

FERTILIZER-FREE ZONES

Section 1-15.07 of the proposed rule states:

"Fertilizer shall not be applied within ten (10) feet from the landward extent of any surface water as identified in section 62-340.600(2), Florida Administrative Code. For example, you may not apply fertilizer within ten feet of the top of bank of any surface water or the top of a seawall."

Justification

Requiring a fertilizer-free and/or "no-mow" zone along water bodies is a standard approach to protecting water quality from impacts due to nutrient enrichment, introduction of oil, grease, and heavy metals, and sediment (Florida Yards & Neighborhoods Handbook, 2009). Keeping impervious surface away from shorelines also reduces the potential for erosion and sedimentation in a water body. Likewise, not applying fertilizer adjacent to surface waters or wetlands greatly decreases the potential for over-enrichment of the water body and subsequent eutrophication. Numerous local governments in Florida now require a setback for site alteration and impervious surfaces, and protection of valuable habitat, within buffer zones (EPC, 2007).

It is widely recognized by professional landscapers and researchers alike that maintaining a non-fertilized strip along water bodies is a good practice for protecting water. No-mow zones also help absorb nutrients present in runoff as well as add a margin for application error (Watschke, 2010; White et al., 2007).

Thus, setting a fertilizer-free and no-mow zone adjacent to water bodies and wetlands is well justified and documented.

Cost / Benefit

Several cost savings can result from implementing a "fertilizer-free zone" adjacent to a surface water body or wetland including the following:

- reduced purchase cost as less overall product would be applied,
- reduced maintenance and labor cost resulting from less area to maintain
- less need to improve degraded water quality through the use of expensive capital improvement projects targeting nutrient-enriched stormwater runoff

OTHER CONSIDERATIONS

Tampa Bay Estuary Program (2006) found that "Stormwater runoff from urban, residential and agricultural lands remains the largest source of nitrogen, the primary pollutant in the bay. An overabundance of nitrogen can cause algae blooms and reduce oxygen levels in the bay, resulting in turbid water, fish kills and loss of seagrass when the water becomes so opaque that sunlight cannot reach underwater grasses. Stormwater accounted for 63% of total nitrogen loadings to Tampa Bay from 1999-2003." More specifically, TBEP found that 20% of the total load to Tampa Bay is estimated from residential stormwater runoff. It should be noted that only a portion of that load is from fertilizers.

The potential benefits of fertilizer management were illustrated in a recent study by the Tampa Bay Estuary Program (2008). TBEP estimates of the potential reduction in N loadings to Tampa Bay were made for ranges of both percent contribution of fertilizer to total urban runoff nutrient loadings, and percent compliance with a series of recommendations for controlling fertilizer application, including a summer blackout with a possible one-time exemption. Conservative estimates from the TBEP indicate an approximate reduction of residential N loading to Tampa Bay by 4.1%.

In addition to the studies and data presented above, other studies indicate the need for a more stringent approach. As stated previously, Leggette, Brashears & Graham, Inc. (LBG) (2004) estimated that roughly 79% of the nitrogen in the surficial groundwater system contributing to Lake Tarpon is from fertilizers (agricultural and residential). Also, a recently published study (2009) by Lehman et al documents statistically significant decreases in total phosphorus concentrations in the Huron River

following implementation of a lawn fertilizer ordinance which restricts the sale of fertilizers containing phosphorus, and requires compliance by lawn care professionals. It should be noted that ambient phosphorus levels in the soils in that part of Michigan largely negate the need for supplemental phosphorus. Nonetheless, this study suggests that a sales restriction can be an effective tool in the management of nutrients into surface water bodies.

Although there are many sources of nutrients that enter surface water and groundwater, studies such as that of Peebles et al (2009) and LBG (2004) indicate that fertilizer use and transport are increasing to an extent that enough enters the receiving water bodies to impact water quality and the local biota.

In addition to these expected water quality benefits, it is anticipated that by limiting a lawn's growth rate it will lower required levels of time and effort for maintenance, as well as provide a cost savings, as recognized by both academic and commercial publications. Trenholm et al. (2006) concluded "*A lower-fertility lawn is best for those with little time to spend on lawn care.... (higher fertility lawns) .. will require more time and money for lawn care.*" Seedland (2009) also stated that "*the level of maintenance you wish to provide (more fertilizer = more work, more mowing more disease control monitoring, and more water) will determine how many applications you make to your grass.*"

SUMMARY and CONCLUSIONS

There are good reasons to adopt an ordinance more stringent than the State's model. Among the most compelling is the fact that multiple water bodies within Hillsborough County, including Tampa Bay, have been declared "impaired waters" by FDEP - with nutrients the causative agents. Impairment is due to nutrient loadings to these water bodies exceeding that which can be assimilated in the water body while maintaining water quality standards and supporting a viable ecosystem. Therefore, actions that reduce the volume of fertilizer applied and which maximize the efficiency of uptake of nutrients by plants, will benefit the environment.

EPC staff asserts that a more stringent rule will provide benefits, including but not limited to:

- prevention of excessive nutrients from entering surface waters due to misuse of products and application techniques
- providing a consistent practice and message throughout the Tampa Bay area since Pinellas County has already passed a similar sales restriction

- allow local governments a cost-effective means of reducing additional nutrient loads to already impaired waters of Hillsborough County
- be a cost-effective mechanism of enforcement and education
- less water required for irrigation, and
- less need to improve degraded water quality through construction of millions of dollars of new and improved stormwater infrastructure projects.

The cost of cleaning polluted water is substantial and local governments usually pay the majority of the amount. Some of the rule provisions discussed above are forms of source control, that is, they reduce the amount of a pollutant before it enters the environment. This is a more economical approach to protecting water quality than remediation, or treating the water downstream after it has become polluted.

Both environmental stewardship and cost savings can be realized while protecting the health and appearance of turf grasses and landscape vegetation by examining established traditional practices and adjusting them to reflect current priorities. The EPC's proposed fertilizer rule, with a vigorous public educational component and manageable enforcement at retail outlets, will be an effective tool to achieve these goals.

Implementation of a source control approach to water quality protection is much less costly than cleaning the water farther downstream after it has become polluted. It has been shown that keeping excess nutrients out of the environment is a superior method of environmental protection in terms of cost to local governments, citizens, and other land owners.

The provisions of the EPC's proposed fertilizer ordinance, including a summer retail sales restriction, increased use of slow-release fertilizer, summer application restrictions, and fertilizer-free zones are necessary as part of an effective, comprehensive approach to controlling nonpoint sources of pollution for the purpose of protecting and improving the aquatic and estuarine resources of Hillsborough County.

Similar to most highly technical and developing fields, opinions about appropriate fertilizer management strategies differ among various entities. However, the discussion above presents credible, technically-defensible evidence that a suite of science-based and economically and technically feasible actions can and should be implemented by the Environmental Protection Commission of Hillsborough County.

REFERENCES

- Alexander, A. and H. Ulrich Helm. 1990. Ureaform as a Slow-release Fertilizer: A Review. In: *Zeitschrift für Pflanzenernährung und Bodenkunde*. V:153, n:4, pp: 249-255.
- American Vegetable Grower. March, 2009. Deliver Better Tubers.
<http://www.growingproduce.com/americanvegetablegrower/index.php?storyid=1462>
- Barth, R. 2009. President, Palmer Ranch Master Property Owners Assoc. Personal Communication.
- Blaylock, A.D., J. Kaufmann, and R.D. Dowbenko. 2005. Nitrogen Fertilizer Technologies. In: *Proceedings, Western Nutrient Management Conference*. 2005. Vol. 6. Salt Lake City, UT.
- Environmental Protection Commission of Hillsborough County. 2007. EPC staff technical findings and recommendations in response to CARE Policy 19.1 of the Comprehensive Plan. Tampa, FL.
- Florida Department of Environmental Protection (FDEP). 2009. 2009 Hillsborough River Basin Management Action Plan for the Implementation of Total Maximum Daily Loads Adopted by the Florida Department of Environmental Protection in the Hillsborough River Basin for Fecal Coliform Bacteria. Tallahassee, Florida.
- Florida Department of Environmental Protection/St. John's River Water Management District (FDEP/SJRWMD). 2010. Final Report Wekiva River Basin Nitrate Sourcing Study. Palatka and Tallahassee, FL.
- Florida Fertilizer Task Force. 2007. Comments on draft State model ordinance. Tallahassee, FL.
- Georgia Pacific. 2009. Nitamin brand fertilizer specifications.
http://www.gp.com/Plant_Nutrition/Default.aspx
- Florida Yards & Neighborhoods Handbook. 2009. Institute of Food and Agricultural Sciences, University of Florida.
- Harmful Algae. December 2008. Volume 8, Issue 1. Edited by P.M. Gilbert, J.M. Burkholder, E. Graneli, and D.M. Anderson. Published by Elsevier. Link: [Science Direct website](#).
- Hauxwell, J., C. Jacoby, T. K. Frazer, J. 2001. Nutrients and Florida's Coastal Waters: The Links Between People, Increased Nutrients, and Changes to Coastal Aquatic Systems. Publication SGEB-55. Florida Sea Grant College Program. Gainesville, FL.
- Hillsborough County Public Works (HCPW). 2009. Personal Communication.
- Hochmuth, G., T. Nell, J. Sartain, B. Unruh, M. Dukes, C. Martinez, L. Trenholm, and J. Cisar. 2009. Unintended Consequences Associated With Certain Urban Turf and Landscape Ordinances. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.
- Hochmuth, G. 2009. UF IFAS. Personal Communication.
- Juchnowicz, M. 2009. President, Gardenmasters of Southwest Florida, Inc. Personal Communication.

- Leggette, Brashears & Graham, Inc. 2004. Lake Tarpon Ground-Water Nutrient Study Final Report prepared for Pinellas County Department of Environmental Management and SWFWMD
- Lehman, J.T., D.W. Bell, K.E. McDonald. 2009. Reduced River Phosphorous Following Implementation of Lawn Fertilizer Ordinance. Lake and Reservoir Management, Vol. 25. University of Michigan.
- National Oceanic and Atmospheric Administration (NOAA). 2010. NOAA National Data Centers (NNDC) Climatological Normals, Normal Precipitation Table for Tampa, Florida.
<http://www.ncdc.noaa.gov/oa/climate/online/ccd/nrmnpp.txt>
- National Oceanic and Atmospheric Administration National Weather Service (NOAA NWS). 2002. The Florida Rain Machine. <http://www.srh.noaa.gov/mlb/?n=flrainmachine>
- Peebles, E., D. Hollander, S. Locker, P. Swarzenski, and G. Brooks. 2009. Areal Extent, Source and Ecological Status of Organic Sediment Accumulation in Safety Harbor, Tampa Bay. Technical Report #07-09 of the Tampa Bay Estuary Program. Prepared by the College of Marine Science, USF.
- Rackley, A. and W. Collier (FDACS). April 2010. Personal Communication.
- Sarasota County Board of County Commissioners. October 26, 2009. Letter from Chair to Pinellas County.
- Sartain, J.B. 2007. General Recommendations for Fertilization of Turfgrasses on Florida Soils. IFAS #SL21. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.
- Sartain, J.B., L.T. Trenholm, E.F. Gilman, T.A. Obreza, and G. Toor. December 2008. Frequently Asked Questions about Landscape Fertilization for Florida-Friendly Landscaping Ordinances. IFAS #ENH1115. Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL.
- Seedland Fertilizers. 2009. Welcome to Lawnfertilizers.com.
<http://www.lawnfertilizers.com>
- Southwest Florida Water Management District. 2002. Annotated Nitrate Bibliography. 147 pages of summaries of studies.
- South Florida Water Management District. 2009 Western Broward County Turf & Landscape Best Management Practices (BMPs).
- Southwest Florida Water Management District. 2009. Fertilizer Facts. Brooksville, FL.
http://www.swfwmd.state.fl.us/yards/fertilizer_facts/details.php#whentoapply
- Tampa Bay Estuary Program, 2006. Charting the Course, the Comprehensive Conservation and Management Plan for Tampa Bay. Published by the Tampa Bay Estuary Program, St. Petersburg, Florida.
- Tampa Bay Estuary Program. 2008. Technical Memorandum: Model-Based Estimates of Nitrogen Load Reductions Associated with Fertilizer Ordinance Implementation. St. Petersburg, FL.
- Thomas, M. (FDEP). April 2010. Personal Communication.
- Trenholm, L.E., J.L. Cisar, and J. Bryan Unruh. 2006. St. Augustine grass for Florida Lawns. Publication ENH5 Institute of Food and Agricultural Sciences. University of Florida. Gainesville, FL.

Troutman, B, E. Santella, J. Lefton, and G. Pickhardt. 2008. Comparative Analysis of Sarasota Model Ordinance Restricted St Augustine Lawns Compared to Traditionally Maintained Lawn in Similar Communities

Wang, F.L. and A.K. Alva. 1996. Leaching of Nitrogen from Slow-Release Urea Sources in Sandy Soils. *Soil Sci Soc Am J* 60:1454-1458

Watschke, T.L. Proper use of fertilizers minimizes environmental effects. *Grounds Maintenance for Golf and Green Industry Professionals*. 2010.

http://www.grounds-mag.com/mag/grounds_maintenance_proper_fertilizers_minimizes/

WeatherBase.com . 2010. Website accessed April 2010. <http://www.weatherbase.com>

White, S.K., H. F. Cook, and L. Garraway. 2007. The Use of Fertilizer-Free Grass Buffer Strips to Attenuate Nitrate Input to Marshland Dykes. *Water and Environment Journal*. Volume 12 Issue 1, Pages 54 - 59

Zone Ten Nursery. 2009. Tree Planting. Miami, FL.

<http://www.zoneten.com/>

Zvomuya, F., Rosen, C.J., Russelle, M.P., and Gupta, S.C. 2003. Nitrate leaching and nitrogen recovery following application of polyolefin-coated urea to potato. *J. Environ. Qual.* 32:480-489.